

Joint Interpretation of Bathymetric and Gravity Anomaly Maps Using Cross and Dot-Products

Pavel Jilinski*

Observatório Nacional, Rio de Janeiro, Brazil

pjilinski@on.br

and

Sergio Luiz Fontes

Observatório Nacional, Rio de Janeiro, Brazil

Summary

We present the results of joint map interpretation technique based on cross and dot-products applied to bathymetric and gravity anomaly gradients maps. According to the theory (Gallardo&Meju, 2004) this technique helps to map unseen relations between bathymetric and gravity anomaly maps when they are analyzed separately and gives information about the correlation between spatial data of different nature.

This method applied for the southern segment of Eastern-Brazilian coast bathymetrical and gravity anomaly gradients indicates a strong source-effect relation between them. The details of the method and the obtained results are discussed.

Introduction

The main objective of this study is to develop and test joint image interpretation method to compare spatial data and analyze its relations.

It is difficult to find correlations between spatial and amplitude distributions of different natural data when one analyzed them separately.

We applied joint map interpretation technique developed by Gallardo and Meju (2004) to investigate the correlation between bathymetric and gravity anomalies at the southern segment of the Eastern-Brazilian coast. Gridded satellite global marine gravity data and bathymetrical data were used.

The studied area is located at the Eastern- Brazilian coast between the 20°W and 30°W meridians and 15°S and 25°S parallels. The volcanic events responsible for the uncommon width of the continental shelf at the Abrolhos bank also were responsible for the formation of the Abrolhos islands and seamounts including the major Vitoria-Trindade chain. According to McKenzie (1976) and Zembruski (1979) such volcanic structures are expected to have a corresponding gravity anomaly.

Theory and Method

Data sources

The bathymetrical satellite data used was derived from bathymetry ETOPO2v2 2-minute grid obtained from NOAA's National Geophysical Data Center (<http://www.ngdc.noaa.gov>).

The satellite marine gravity 1-minute gridded data were obtained from the Satellite Geodesy at the Scripps Institution of Oceanography, Smith & Sandwell (1997). (<http://topex.ucsd.edu>).

Gravity anomaly data were re-gridded using the ETOPO2v2 grid.

Cross-Product

Cross-product is the result of multiplication of bathymetric and gravity anomaly gradient magnitudes by the sine of the angle between them.

According to the definition of gradient cross-product, minimal values are expected to be found in points where the angle between gradients is close to zero or where one or both of the gradient magnitudes have values close to zero. It creates an ambiguity and problems for data interpretation since there is no exact correspondence between bathymetric structures and gravity anomalies.

Dot-Product

Dot-product is the result of multiplication of bathymetric and gravity anomaly magnitudes by the cosine on the angle between them.

According to the definition of dot-product, minimal values can be generated by near perpendicular orientation of the gradients or small magnitudes of one or both gradients. So, the results are mutually increased in areas with larger magnitudes or smaller angles between gradients. Due to this mutual amplification dot-products are less affected by the explained above ambiguity of cross-product. Some statistical treatments of cross-product were used to support the conclusions.

Statistics and Significance Criteria

Statistical analysis was made in order to sort the magnitude data into two groups to reduce ambiguity effect: first group – data with magnitudes that could be considered *anomalous* (where the main minimizing source is the angle between the gradients) and the second group – data with magnitudes variations that could be considered as non significant or *background* (where cross-product value is determined by the small magnitude). It was chosen to use the mean value and standard deviation to sort the data in such two groups. Those values were determined for bathymetric and gravity anomaly gradient magnitudes creating two data sets – one where one or both gradient magnitudes are one standard deviation larger than the mean value with a total number of 7831 (*anomalous*) points and a second one where both magnitudes differ smaller than one standard deviation from the mean value with total number of 85584 (*background*) points. Statistical analysis of distribution patterns for both groups was made.

Examples of Method Application

Map of Angles Between Gradients

The angle values were divided into 4 equal intervals. The statistical distribution of angles between gradient in the given intervals is the following (percents of the total): 0 to 60° – 51.39% of the values; 60° to 90° – 12.08%; 90° to 120° – 14.92%; 120° to 180° – 21.18%.

It can be seen that 51% of the gradients have a small angle between them, 72% of gradients can be considered as "parallel" with angles smaller than 60° or bigger than 120° between them.

After statistical separation in the *anomalous* group almost 91% of the gradients have an angle smaller than 60° while in the *background* group this value is just 48.6%.

From these results we can make a conclusion that the majority of the bathymetric and gravity anomaly gradients are related.

Regions with higher gradient magnitudes are characterized by cosine values close to 1 (indicating a small angle between them). The areas characterized by small angles between gradients exceed the size of bathymetric structures and gravity anomaly characterizing the influence of the structures and their effects on surrounding areas.

Regions with no significant anomalies show uncorrelated value spots. Figure 1 shows the map of angle values between bathymetric and gravity gradients.




Figure 1: Map of Angles Between Bathymetric and Gravity Anomaly Gradients for the Studied Region

Map of Cross-Product

The resulting map (Figure 2 (a)) shows small spots of higher cross-product magnitudes following magnitude isolines. About 90% of the values are close to minimum.




Figure 2: Cross product (a) and Dot product (b) maps of the same region as Fig. 1.

As was mentioned above, we can presume that areas where bathymetry and gravity anomaly gradient cross-products have smallest magnitudes there is a good correspondence between them indicating a good correspondence between shapes.

According to these results for the studied area the shapes and positions of bathymetric structures and gravity anomalies are well correlated suggesting strong correlation between source and its effect.

Map of Dot-Product

The dot-product map of the studied area (Fig. 2 (b)) resembles bathymetric and gravity anomaly isolines and suggests a strong source-effect between bathymetry and gravity anomaly.

All the sea mounts, banks, continental slope and other notable geomorphologic structures and gravity anomalies are well delimited in the dot-product map eliminating uncorrelated areas where gradient orientations can be considered as near perpendicular.

Conclusions

It is shown that the joint image interpretation technique uses three different criteria which are sensitive to different gradient properties.

Angles between gradients are a good indicator of areas where data are related and it is not sensitive to the magnitudes of the gradients. Angles maps can be used to find areas with direct and inverse relation between mapped properties and contour areas of influence of anomalies unseen on gradient magnitude maps alone. Statistical measures of distribution of angles can be an indicator of relations between two data sets as show using significance criteria.

Cross-product map has a *spotted* character of contours. To reduce the effects of the ambiguity the separation of the data into two groups proved to be useful. It helped to separate the cross-product values that are minimized due to gradient magnitudes from those that minimize due to sine values which is a measure of correlation between them.

Dot-product values contour areas where gradients are correlated.

According to joint image interpretation technique applied bathymetric structures especially the volcanic seamounts and banks in the southern part of East-Brazilian Coast are closely related to the observed gravity anomalies and can be interpreted as sources and effect. This technique also helps to evaluate the shape and dispersion of the gravitational effect from a bathymetrical source.

References

- Dehlinger P., Marine Gravity, Elsevier, 1978.
- Gallardo, L. A., and M. A. Meju., Joint 2D cross-gradient imaging of magnetotelluric and seismic travel-time data for structural and lithological classification, *Geophys. J. Int.*, 169, 1261-1272. (2007)
- Gallardo, L.A., M. A. Meju (2004), Joint two-dimensional dc resistivity and seismic travelttime inversion with cross-gradients constraints, *J. Geophys. Res.*, 109, B03311, doi:10.1029/2003JB002716
- Jacoby, W., and Smilde P. L., Gravity Interpretation, Springer, 2009.
- McKenzie D. & Bowin C. 1976. The relationship between bathymetry and gravity in Atlantic Ocean. *Journal of Geophysical Research*, 81: 1903-1915.
- Roy. K. K., Potential Theory in Applied Geophysics, Springer, 2008.
- Smith, W. H. F., and D. T. Sandwell, Global seafloor topography from satellite altimetry and ship depth soundings, *Science*, v. 277, p. 1957-1962, 26 Sept., 1997.
- Sandwell, D. T., and W. H. F. Smith, Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge Segmentation versus spreading rate, *J. Geophys. Res.*, 114, B01411, doi:10.1029/2008JB006008, 2009.
- Zembruscki, S.G. 1979. Geomorfologia da Margem Continental Sul Brasileira e das Bacias Oceânicas Adjacentes. *In: Geomorfologia da margem continental brasileira e das áreas oceânicas adjacentes. Série Projeto REMAC, N° 7.*